

SEVERE ACUTE RESPIRATORY SYNDROME

This application claims priority from U.S. Provisional Application No. 60/468,644, filed May 8, 2003, the entire content of which is incorporated
5 herein by reference.

TECHNICAL FIELD

The present invention relates, in general, to severe acute respiratory syndrome (SARS) and, in particular, to a method of generating neutralizing
10 antibodies to the virus. The invention further relates to a method of detecting the presence of the virus and to a method of treating an infected individual.

BACKGROUND

15 Since the severe acute respiratory syndrome (SARS) epidemic surfaced in Asia, more than 2600 cases have been identified in 19 countries, and more than 100 deaths have been reported. SARS has recently been identified as a new clinical entity
20 (INFECTIOUS DISEASES: Deferring Competition, Global Net Closes In on SARS. Science 300(5617):224-5 (2003); Ksiazek et al, N. Engl. J. Med. Apr 10 (2003); Drosten et al, N. Engl. J. Med. Apr 10 [epub ahead of print] (2003); Poutanen et al, N. Engl. J.
25 Med. Apr 10 [epub ahead of print] (2003)). It has been found that a novel coronavirus is associated with this outbreak, and the evidence indicates that this virus has an etiologic role in SARS since this

virus was found in samples from multiple SARS patients in several independent laboratories. The complete genome of the SARS associated coronavirus ("the SARS virus") was derived by sequencing of gene
5 fragments generated using consensus coronavirus primers designed to amplify SARS genes by reverse transcription-polymerase chain reaction (RT-PCR).

The SARS virus is RNA virus with the genome size of approximately 29K nucleotides. The complete
10 SARS virus genome sequence has been reported by Jones et al and is available in the NCBI DNA database (GI: 29826277). Phylogenetic analyses and sequence comparisons showed that the SARS virus is not closely related to any of the previously
15 characterized coronaviruses (Figs. 1-5).

SUMMARY OF THE INVENTION

The present invention relates generally to SARS. More specifically, the invention relates to a method of producing neutralizing antibodies to the
20 virus and to a method of treating individuals infected with the virus. The invention further relates to a method of detecting the presence of the virus in a sample. The invention additionally relates to compounds and compositions suitable for
25 use in such methods.

Objects and advantages of the present invention will be clear from the description that follows.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Amino acid sequence comparison of spike protein between SARS coronavirus with bovine coronavirus.

5 Figure 2. Amino acid sequence comparison of spike proteins between SARS coronavirus with human coronavirus OC43.

Figure 3. Phylogenetic analysis of coronavirus N protein.

10 Figure 4. Phylogenetic analysis of coronavirus S protein.

Figure 5. Phylogenetic analysis of coronavirus M protein.

15 Figure 6. Protein structure of SARS virus spike glycoprotein.

Figure 7. Protein structure of SARS virus nucleocapsid (NP) protein.

Figure 8. SARS spike protein peptides.

Figure 9. SARS NP protein peptides.

Figure 10. Coronavirus spike protein among isolates.

Figure 11. Peptide design based on predicated SARS spike protein antigenic epitopes.

5 Figure 12. HR and LZ domains in coronavirus spike proteins. (HR1 (SEQ ID NO:34), HR2 (SEQ ID NO:35))

Figure 13. Immunization protocol of rabbits with SARS spike protein peptides.

10 Figure 14. Schematic representation of SARS expression vectors.

Figure 15. Western blot analysis of SARS spike protein, shown are purified SARS spike protein (lane 1), spike protein Ig fusion protein (lane 3) and mock transfection supernatant control, produced in transformed 293 cells and purified using a lectin column - analysis was effected using Western blot and detection using immune sera of a mouse immunized with a DNA vaccine expressing SARS spike protein.

15

20 Figure 16. Induction of antibody reacted with recombinant SARS spike protein by immunization with plasmid DNAs that express SARS-spike protein or spike protein-Ig. Serum samples were collected

10 days after immunizations and assayed by ELISA.
Shown are the end-point ELISA titers against
recombinant SARS spike proteins coated on a 96-well
plate (200 ng/well).

5

DETAILED DESCRIPTION OF THE INVENTION

In one embodiment, the present invention
relates to a method of producing neutralizing
antibodies to the SARS virus. In a further
embodiment, the invention relates to a method of
10 treating an individual infected with the virus. In
another embodiment, the invention relates to a
method of detecting the presence of the SARS virus
in a sample (e.g.. a biological sample). The
invention also relates to compounds and compositions
15 suitable for use in the such methods.

The structure of the SARS virus putative spike
glycoprotein (1,255 amino acids) and that of the
nucleocapsid protein (NP) (422 amino acids) have
been analyzed using DNASTar computer program,
20 version 3.16 (DNASTar Inc.) (see Figs. 6 and 7,
respectively; the notation on the right margin
indicates the nature of the region such as
antigenicity index, surface probability etc.).

Based on the antigenic index of these two
25 proteins, and data in the literature relating to
other coronaviruses, the panel of peptides listed in
Table 1 (SEQ ID NO:1 to SEQ ID NO:33, respectively)
has been designed (see also Fig. 8 and 9).
Positions of variability that have been identified

in the SARS virus spike protein are shown in Fig.
10.

Table 1. Synthetic Peptides derived from SARS coronavirus spike and N proteins.

Name of peptide	Amino acid sequence	a.a position
DUHVI SA-S1	TTFDDVQAPNYTQHTSSMRGVYYPDEIFRSDT	20-51 *
DUHVI SA-S2	FKDGIYFAATEKSNVVRGWVFGSTMNKSQS	83-113
DUHVI SA-S3	NSTNVVIRACNFELCDNPFFAVSKPMGTQTH	119-149
DUHVI SA-S4-A	FEYISDAFSLDVSEKSGNFKHLREFVFK	161-188 *
DUHVI SA-S4	DVSEKSGNFKHLREFVFKNKDGFYLVYKGYQPIDVVRDLPS	171-213
DUHVI SA-S4-B	KGYQPIDVVRDLPSGFNTLKPIFK	198-221 *
DUHVI SA-S5	FSPAQDIWGTSAAYFVGYLKPTTFMLKYDENGTTIT	238-273
DUHVI SA-S6	KYDENGTTITDAVDCSQNPLAELK	265-287 *
DUHVI SA-S7	FSPAQDIWGTSAAYFVGYLKPTTFMLKYDENGTTIT	288-320
DUHVI SA-S8	FVVGDDVRQIAPGQTGVYADVNYKLPDDFM	386-417
DUHVI SA-S9	NTRNIDATSTGNYNKYRYLRHGKLRPFERDISN	424-457 *
DUHVI SA-S10	FSPDGKPCPTPALNCYWPLNDYGFYTTTGIG	460-490
DUHVI SA-S11	PKLSTDLIKNCQVNFNFNGLTGTGVLTPSSKRFQ	513-546
DUHVI SA-S12	TPSSKRFQPFQFGRDVSDFDTSVRDPKTSE	539-569 *
DUHVI SA-S13	TNASSEVAVLYQDVNCTDVSTAIHADQLTPAWRIYSTGN	588-626
DUHVI SA-S14	EHVDTSYECDIPIGAGICASYHTVSLLRSTSQKSI	640-674
DUHVI SA-S15	EHVDTSYECDIPIGAGICASYHTVSLLRSTSQKSI	753-782
DUHVI SA-S16	LKPTKRSFIEDLLFNKVTLAGAFMKQYGECLGDINARDL	792-831
DUHVI SA-S17	NQKQIANQFNKAISQIQESLTTTSTALGKLQDVVNQNAQ	901-939
DUHVI SA-S18	SKRVDFCGKGYHLMSFPQAAPHGVVFLHVTYVPSQERNF	1019-1057
DUHVI SA-S19	EGKAYFPREGVVFVNGTSWFITQRNFFSP	1066-1094
DUHVI SA-S20	DPLQPELDSFKEELDKYFKNHTSPDVLGDISG	1121-1153 *
DUHVI SA-S21	QKEIDRLNEVAKNLNESLIDLQELGKYEQY	1162-1191
DUHVI SA-S22	LTVLPPLLTDDMIAAYTAALVSGTATAGWTFGAGAALQIPF	841-882 *
DUHVI SA-S23	AMQMAYRFNGIGVTONVLYENQKQIANQFNKAISQIQESL	843-921 *
DUHVI SA-S24	ELDSFKEELDKYFKNHTSPDVLGDISGINASVV	1127-1161 *
DUHVI SA-S25	NIQKEIDRLNEVAKNLNESLIDLQELGKYEQYIKWPW	1162-1197 *
DHVI SA-N1	DSTDNNQNGGRNGARPKQRRPQGLENN	23-49 *
DHVI SA-N2	GSRGGSQASSRSSRSRSGNSRNSTPGSSRGNSPAR	176-210 *
DHVI SA-N3	KVSGKGQQQQGQTVTKKSAEASKKPRQKRTATK	234-267 *
DHVI SA-N4	GRRGPEQTQGNFGDQDLIRQGTDYKH	276-301 *
DHVI SA-N5	HIDAYKTFFPTEPKDKKKKTDEAQPLPQRQKKQ	357-369
DHVI SA-N6	QKKQPTVTLLPAADMDDFSRQLQNSMSGASADSTQ	387-421

5 The present invention includes the peptides set forth in Table 1 (and Figs. 8 and 9), corresponding peptides from other SARS virus isolates and unique and/or antigenic portions of such peptides. Unique and/or antigenic portions are preferably at least 5

amino acids in length, more preferably, at least 6, 7, 8, 9 or 10 amino acids in length. The peptides can be synthesized, for example, using standard chemical syntheses techniques, as can polymers
5 containing multiple copies of one or more of the above peptides or portions. The peptides (portions and polymers) can also be synthesized using well-known recombinant DNA techniques. Recombinant synthesis may be preferred when the peptides are
10 covalently linked.

In addition to the above peptides (and portions and polymers), the invention also relates to nucleic acids encoding the same. The nucleic acids (e.g., DNA) can be present in a vector (e.g., a viral
15 vector or a plasmid), advantageously linked to a promoter.

The invention includes compositions containing one or more of the above peptides (or portions or polymers), or nucleic acids encoding same, and a
20 carrier, e.g., a pharmaceutically acceptable carrier. The peptide-containing compositions can further include an adjuvant (such as alum). The peptides of the invention (or portions or polymers) can be present in the composition conjugated to a
25 carrier molecule, either directly or indirectly via a spacer molecule. Carrier molecules are, advantageously, non-toxic, pharmaceutically acceptable and of a size sufficient to produce an immune response in mammals. Examples of suitable
30 carriers include tetanus toxoid and keyhole limpet hemocyanin.

As indicated above, in one embodiment, the present invention relates to a method of producing neutralizing antibodies in a mammal (e.g., a human) to the SARS virus. The method comprises

5 administering to a mammal in need thereof an amount of one or more of the above-described peptides, portions or polymers, sufficient to effect the production of neutralizing antibodies. (See also Figs. 11 and 12 - the regions specifically depicted

10 in Fig. 11 corresponding to regions reportedly associated with the induction of neutralizing antibodies in the context of other coronaviruses; Fig. 12 provides the sequences of HR1 and HR2 - these are sequences demonstrated to be capable of

15 inhibiting fusion of animal coronaviruses (see Daniel et al, J. Virol. 67:1185-1194 (1993); Routledge et al, J. Virol. 65:254-262 (1991); Talbot et al. J. Virol 62:3032-3036 (1988) and Luo and Weiss In Coronavirus and Arteriviruses, ed. by

20 Enjuanes, pp. 17-22 (1998)).) Optimum dosing regimens, which can vary with the peptide used, the patient and the effect sought, can be readily determined by one skilled in the art.

In an alternative aspect of this embodiment,

25 production of neutralizing antibodies to the SARS virus can be effected by administering the above-described nucleic acids under conditions such that the nucleic acid is expressed, the encoded peptide produced and the neutralizing antibodies generated.

30 That is, nucleic acids encoding the peptides (portions and polymers) of the invention can be used

as components of, for example, a DNA vaccine wherein the peptide encoding sequence(s) is/are administered as naked DNA or, for example, a minigene encoding the peptides can be present in a viral vector. The
5 encoding sequence(s) can be present, for example, in a replicating or non-replicating adenoviral vector, an adeno-associated virus vector, an attenuated mycobacterium tuberculosis vector, a Bacillus Calmette Guerin (BCG) vector, a vaccinia or Modified
10 Vaccinia Ankara (MVA) vector, another pox virus vector, recombinant polio and other enteric virus vector, Salmonella species bacterial vector, Shigella species bacterial vector, Venezuelan Equine Encephalitis Virus (VEE) vector, a Semliki
15 Forest Virus vector, or a Tobacco Mosaic Virus vector. The encoding sequence(s), can also be expressed as a DNA plasmid with, for example, an active promoter such as a CMV promoter. Other live vectors can also be used to express the sequences of
20 the invention. Expression of the peptides of the invention can be induced in a patient's own cells, by introduction into those cells of nucleic acids that encode the peptides, preferably using codons and promoters that optimize expression in human
25 cells. Examples of methods of making and using DNA vaccines are disclosed in U.S. Pat. Nos. 5,580,859, 5,589,466, and 5,703,055.

In another embodiment, the present invention relates to a method of treating an individual (e.g.,
30 a human) infected with the SARS virus. As above, this method can be effected by administering the

above-described peptides (portions and polymers)
(the use of one or more of peptides SA-20 to SA-25
from Table 1, or portions thereof or polymers
comprising same, being preferred) or nucleic acids
5 in an amount and under conditions such that the
treatment is effected. Peptides comprising HR-1
and/or HR-2, or portions thereof, are particularly
preferred. The significance of the HR-1 and HR-2
(LZ (leucine zipper)) regions is that these are
10 homologous regions to the coil coil structures of
HIV gp41, and HR-2 corresponds to the HR-2 or (T-20)
drug that is working so well for HIV. Thus, the
SARS virus HR-1 or HR-2 peptide (or portion thereof)
can be expected to inhibit fusion of infected cells
15 and prevent virus entry.

Optimum dosing regimens can be readily
determined by one skilled in the art.

Suitable routes of administration of the
peptides (portions and polymers) and nucleic acid of
20 the invention include systemic (e.g. intramuscular
or subcutaneous). Alternative routes can be used
when an immune response is sought in a mucosal
immune system (e.g., intranasal).

In another embodiment, the invention relates to
25 methods of detecting the SARS virus in a sample
(e.g., a biological sample from a patient, such as a
blood, serum, sputum or fecal sample, or an
environmental sample, such as a water or sewage
sample). As appropriate, the method can be effected
30 by detecting the presence of viral proteins or
nucleic acids. For example, the above-described

peptides (portions or polymers) can be used to generate antibodies (polyclonal or monoclonal) using standard techniques. The antibodies (or binding fragments thereof) can then be used, for example, in
5 standard immunoassays, to detect the presence of SARS viral protein in the sample. The peptides (portions and polymers) can also be used, for example, in accordance with standard immunoassay techniques, to detect the presence of viral
10 antibodies in, for example, the blood of a patient. Alternatively, the nucleic acids described above, or complements thereof, can be used according to standard techniques as probes or primers to detect the presence of viral encoding sequences in a
15 sample. It will be appreciated that any of the peptides (portions or polymers), antibodies (or fragment) or nucleic acids can bear a detectable label (e.g., a fluorescent or radiolabel).

Certain aspects of the invention can be
20 described in greater detail in the non-limiting Examples that follows.

EXAMPLE 1

Development of polyclonal immune sera by
immunization in rabbits with synthetic peptides
25 derived from SARS virus

Peptides listed in Table 1 are synthesized as crude peptides, purified and analyzed. Rabbits (2 for each peptides) are immunized with this panel of

SARS virus peptides at a dose of 250µg per injection per animal for a total of 5 immunizations with RIBI adjuvant. Serum samples are collected 10 days after each immunization, and assayed against the
5 immunizing peptides. Further characterization of immune sera including the reactivity of immune sera with native SARS virus proteins is effected.

EXAMPLE 2

10

Development of monoclonal antibodies against the SARS virus spike glycoprotein and NP using synthetic peptides derived the SARS virus as immunogen

15 Based on the initial immunogenicity results of the panel of SARS virus peptides, 1-2 peptides are selected from both SARS spike glycoprotein and NP as immunogens to immunize Balb/c mice for development of monoclonal antibodies. Immune sera and initial
20 screening of hybridoma cell culture are carried out using the immunizing peptides. Further characterization and screening of monoclonal antibodies are effected using SARS native spike glycoprotein and NP expressed in a eukaryotic cell
25 expression system. The neutralizing activities of the monoclonal antibodies are assessed.

EXAMPLE 3

Development of polyclonal immune sera by
immunization of rabbits with synthetic peptides
5 derived from SARS coronavirus.

The protein structure of the putative spike glycoprotein (1,255 amino acids) has been analyzed using DNASTar computer program. Based on the
10 antigenic index of these two proteins, a panel of 33 peptides derived from SARS coronavirus spike protein and NP proteins (as listed in Table 1) has been designed. Of these peptides, nine (S1, S4A, S4B, S9, S12, S20, S23, S24 and S25) have been used to
15 immunize rabbits using a immunization protocol as shown in Figure 13. Other peptides will be used in the future experiments.

EXAMPLE 4

20 Expression of SARS coronavirus spike glycoprotein
and development of monoclonal antibodies (Mabs)
against SARS virus.

25 To develop Mabs and vaccine immunogens against SARS virus, a SARS coronavirus spike protein gene has been developed with codon- and RNA structure optimized for optimal expression. To produce secreted soluble SARS spike protein, an expression
30 vector (SARS SΔTC) was generated in which the transmembrane (TM) and cytoplasmic domain (Cyt) of

SARS spike protein was deleted. To enhance the immunogenicity and stability as well as to provide for ease of purification of SARS spike protein, the extracellular domain of SARS spike protein was
5 linked with either mouse or human IgG constant region genomic sequence (Figure 14). These 2 vectors were used for production of spike protein *in vitro* by transfection and also used as vaccine immunogens for development of monoclonal antibody as well as
10 vaccine immunogens for induction of neutralizing antibodies against SARS virus.

As shown in Figure 15, SARS spike proteins have been expressed in 293 cells by transfection with SARS SATC and SARSATC-Ig vectors and purified using
15 a lectin column. Purified proteins were analyzed by SDS-PAGE and Western blot (Figure 15). The extracellular domain SARS spike protein has a molecular weight of approximately 150Kda, and SARS spike protein-Ig fusion protein has a molecular
20 weight of approximately 170Kda as detected by immune serum from a mouse immunized with the DNA vaccine that expresses SARS spike protein extracellular domain (Figure 14). The purified SARS spike protein has been used for evaluation of immunogenicity of
25 SARS spike protein expression DNA vaccine (see below). To generate Mabs, mice (4 mice for each group) have been immunized with the SARS SATC vector that expresses SARS spike protein. Mice developed antibody responses as detected using Western blot
30 (Figure 15) and ELISA (Figure 16). Both SARS SATC and SARSATC-Ig vectors were also used as DNA vaccine

immunogens for evaluation of the immunogenicity for
induction of neutralizing antibody against SARS.

* * *

All documents cited above are hereby
5 incorporated in their entirety by reference.